

Role of the singular factors in the standard fits for initial parton densities

B.I. Ermolaev

Ioffe Physico-Technical Institute, 194021 St.Petersburg, Russia

M. Greco

Department of Physics and INFN, University Rome III, Rome, Italy

S.I. Troyan

St.Petersburg Institute of Nuclear Physics, 188300 Gatchina, Russia

Total resummation of double- and single- logarithms of x contributing to the spin-dependent structure function g_1 ensures its steep rise at small x . In the asymptotic limit $x \rightarrow 0$, the resummation leads to the Regge behavior of g_1 and allows to calculate the non-singlet and singlet intercepts of g_1 . DGLAP lacks such a resummation but suggests special phenomenological fits for the initial parton densities such that the singular factors $x^{-\alpha}$ in the fits mimic the resummation and also provide g_1 with the steep (power-like) rise at the small- x region. Accounting for the total resummation of logarithms of x allows to drop the singular factors in the fits and leads to a remarkable simplification of the fits.

PACS numbers: 12.38.Cy

I. INTRODUCTION

The standard theoretical instrument for investigating the DIS structure function $g_1(x, Q^2)$ is DGLAP[1]. In this approach, g_1^{DGLAP} is a convolution of the coefficient functions C_{DGLAP} and evolved parton distributions which are also expressed as a convolution of the splitting functions P_{DGLAP} and initial parton densities. The latter are found with fitting experimental data at x close 1 and $Q^2 \sim 1 \text{ GeV}^2$. There is an obvious asymmetry in treating Q^2 - and x - contributions in DGLAP. Indeed, the leading Q^2 - contributions, $\ln(Q^2)$, are accounted to all orders in α_s whereas $C_{DGLAP}(x)$ and $P_{DGLAP}(x)$ are known in first two orders of the perturbative QCD. The reason for such asymmetry is the fact that originally DGLAP was constructed for operating at large Q^2 and x not so far from 1. In this region $\ln^k Q^2$ were large whereas x - contributions from higher loops were small and could be neglected. On the contrary, in the small- x region the situation looks opposite: logarithms of x , namely double logarithmic (DL), i.e. the terms $(\alpha_s \ln^2(1/x))^k$, and single logarithms (SL), the terms $(\alpha_s \ln(1/x))^k$, with $k = 1, 2, \dots$, are becoming quite sizable and should be accounted to all orders in α_s . When the total resummation of DL terms was done[2], it led to new expressions, g_1^{DL} , for g_1 . In particular, Refs. [2] demonstrated that the small- x asymptotics of g_1^{DL} was of the Regge (power-like) form and was much greater than the well-known small- x asymptotics of g_1^{DGLAP} (see Eq. (3)) at $x \leq 0.01$. However, this result was strongly criticized (see e.g. Refs. [3]) where it was shown that using the standard NLO DGLAP formulae together with appropriate fits for initial parton densities led to the opposite result: $g_1^{DL} \ll g_1^{DGLAP}$ at small x . After that the total resummation of logarithms of x for g_1 was claimed irrelevant for available range of x .

Strictly speaking, comparison of results of Refs. [2] and [3] could not be done in a straightforward way because DGLAP -evolution equations have always used the running α_s , with the parametrization

$$\alpha_s^{DGLAP} = \alpha_s(Q^2), \quad (1)$$

whereas Refs. [2] operated with α_s fixed at an unknown scale. Trying to set a scale for α_s when the approach of Refs. [2] is used, Refs. [4] suggested that the argument of α_s should be Q^2 like in DGLAP. The parametrization $\alpha_s^{DGLAP} = \alpha_s(Q^2)$ appears when the argument of α_s in each of the ladder rungs of the involved Feynman graphs is $k_{\perp}^2$, with k being the momentum of the upper parton (a quark or a gluon) of the rung. A deeper investigation of this matter[5] led us to the conclusion that the DGLAP- parametrization of Eq. (1) can be a good approximation at x not far from 1 only. Instead of it, a new parametrization was suggested where the argument of α_s in every ladder rungs is the virtuality of the horizontal gluon (see Ref. [5] for detail). This parametrization is universally good for both small x and large x . It converges to the DGLAP- parametrization at large x but differs from it at small x . Using this new parametrization allowed us to obtain in Refs. [7] expressions for g_1 accounting for all-order resummations of DL and

SL terms, including the running α_s effects¹. In the first place, it was used to obtain numerical values of the intercepts of the singlet and non-singlet g_1 . It is worth to mention that these results were immediately confirmed[8] by several independent groups who fitted available HERMES data and extrapolated them into the asymptotic region $x \rightarrow 0$.

Nevertheless, it is known that, despite DGLAP lacks the total resummation of $\ln x$, it successfully operates at $x \ll 1$. As a result, the common opinion was formed that not only the total resummation of DL contributions in Refs. [2] but also the much more accurate calculations performed in Refs. [7] should be out of use at available x and might be of some importance in a distant future at extremely small x . In Ref. [9] we argued against such a point of view and explained why DGLAP can be so successful at small x : in order to be able to describe the available experimental data, DGLAP uses the singular fits (see for example Refs. [10, 11]) for the initial parton densities. Singular factors (i.e. the factors which $\rightarrow \infty$ when $x \rightarrow 0$) in the fits mimic the total resummation of Refs. [7]. Using the results of Ref [7] allows to simplify the quite complicated structure of the standard DGLAP fits down to normalization constants at small x .

II. DIFFERENCE BETWEEN DGLAP AND OUR APPROACH

In DGLAP, g_1 is expressed through convolutions of the coefficient functions and evolved parton distributions. As convolutions look simpler in terms of integral transforms, it is convenient to represent g_1 in the form of the Mellin integral. For example, the non-singlet component of g_1 can be represented as follows:

$$g_1^{NS}{}_{DGLAP}(x, Q^2) = (e_q^2/2) \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} (1/x)^\omega C_{DGLAP}(\omega) \delta q(\omega) \exp \left[\int_{\mu^2}^{Q^2} \frac{dk_\perp^2}{k_\perp^2} \gamma_{DGLAP}(\omega, \alpha_s(k_\perp^2)) \right] \quad (2)$$

with $C_{DGLAP}(\omega)$ being the non-singlet coefficient functions, $\gamma_{DGLAP}(\omega, \alpha_s)$ the non-singlet anomalous dimensions and $\delta q(\omega)$ the initial non-singlet quark densities in the Mellin (momentum) space. The expression for the singlet g_1 is similar, though more involved. Both γ_{DGLAP} and C_{DGLAP} are known in first two orders of the perturbative QCD. Technically, it is simpler to calculate them at integer values of $\omega = n$. In this case, the integrand of Eq. (2) is called the n -th momentum of g_1^{NS} . When the moments for different n are known, g^{NS} at arbitrary values of ω is obtained with interpolation of the moments. Expressions for the initial quark densities are defined from phenomenological consideration, with fitting experimental data at $x \sim 1$. Eq. (2) shows that γ_{DGLAP} govern the Q^2 - evolution whereas C_{DGLAP} evolve $\delta q(\omega)$ in the x -space from $x \sim 1$ into the small x region. When, at the x -space, the initial parton distributions $\delta q(x)$ are regular in x , i.e. do not $\rightarrow \infty$ when $x \rightarrow 0$, the small- x asymptotics of $g_1{}_{DGLAP}$ is given by the well-known expression:

$$g_1^{NS}{}_{DGLAP}, \quad g_1^S{}_{DGLAP} \sim \exp \left[\sqrt{\ln(1/x) \ln \left(\ln(Q^2/\mu^2) / \ln(\mu^2/\Lambda_{QCD}^2) \right)} \right]. \quad (3)$$

On the contrary, when the total resummation of the double-logarithms (DL) and single- logarithms of x is done[5], the Mellin representation for g_1^{NS} is

$$g_1^{NS}(x, Q^2) = (e_q^2/2) \int_{-\infty}^{\infty} \frac{d\omega}{2\pi i} (1/x)^\omega C_{NS}(\omega) \delta q(\omega) \exp \left(H_{NS}(\omega) \ln(Q^2/\mu^2) \right), \quad (4)$$

with new coefficient functions C_{NS} ,

$$C_{NS}(\omega) = \frac{\omega}{\omega - H_{NS}(\omega)} \quad (5)$$

and anomalous dimensions H_{NS} ,

$$H_{NS} = (1/2) \left[\omega - \sqrt{\omega^2 - B(\omega)} \right] \quad (6)$$

where

$$B(\omega) = (4\pi C_F(1 + \omega/2)A(\omega) + D(\omega))/(2\pi^2). \quad (7)$$

¹ The parametrization of see Ref. [5] was used later in Refs. [6] for studying the small- x contribution to the Bjorken sum rule.

$D(\omega)$ and $A(\omega)$ in Eq. (7) are expressed in terms of $\rho = \ln(1/x)$, $\eta = \ln(\mu^2/\Lambda_{QCD}^2)$, $b = (33 - 2n_f)/12\pi$ and the color factors $C_F = 4/3$, $N = 3$:

$$D(\omega) = \frac{2C_F}{b^2N} \int_0^\infty d\rho e^{-\omega\rho} \ln\left(\frac{\rho+\eta}{\eta}\right) \left[\frac{\rho+\eta}{(\rho+\eta)^2 + \pi^2} + \frac{1}{\rho+\eta} \right], \quad (8)$$

$$A(\omega) = \frac{1}{b} \left[\frac{\eta}{\eta^2 + \pi^2} - \int_0^\infty \frac{d\rho e^{-\omega\rho}}{(\rho+\eta)^2 + \pi^2} \right]. \quad (9)$$

H_S and C_{NS} account for DL and SL contributions to all orders in α_s . When $x \rightarrow 0$,

$$g_1^{NS} \sim (x^2/Q^2)^{\Delta_{NS}/2}, \quad g_1^S \sim (x^2/Q^2)^{\Delta_S/2} \quad (10)$$

where the non-singlet and singlet intercepts are $\Delta_{NS} = 0.42$, $\Delta_S = 0.86$. The x -behavior of Eq. (10) is much steeper than the one of Eq. (3). Obviously, the total resummation of logarithms of x leads to the faster growth of g_1 when x decreasing compared to the one predicted by DGLAP, providing the input δq in Eq. (2) is a regular function of ω at $\omega \rightarrow 0$.

III. STRUCTURE OF THE STANDARD DGLAP FITS

Although there are different fits for $\delta q(x)$ in literature, all available fits include both regular and singular factors when $x \rightarrow 0$ (see e.g. Refs. [10, 11] for detail). For example, one of the fits from Ref. [10] is

$$\delta q(x) = N\eta x^{-\alpha} \left[(1-x)^\beta (1+\gamma x^\delta) \right], \quad (11)$$

with N , η being a normalization, $\alpha = 0.576$, $\beta = 2.67$, $\gamma = 34.36$ and $\delta = 0.75$. In the ω -space Eq. (11) is a sum of pole contributions:

$$\delta q(\omega) = N\eta \left[(\omega - \alpha)^{-1} + \sum m_k (\omega + \lambda_k)^{-1} \right], \quad (12)$$

with $\lambda_k > 0$, so that the first term in Eq. (12) corresponds to the singular factor $x^{-\alpha}$ of Eq. (11). When the fit Eq. (11) is substituted in Eq. (2), the singular factor $x^{-\alpha}$ affects the small- x behavior of g_1 and changes its asymptotics Eq. (3) for g_1 for the Regge asymptotics. Indeed, the small- x asymptotics is governed by the leading singularity $\omega = \alpha$, so

$$g_1^{DGLAP} \sim C(\alpha) (1/x)^\alpha \left((\ln(Q^2/\Lambda^2))/(\ln(\mu^2/\Lambda^2)) \right)^{\gamma(\alpha)}. \quad (13)$$

Obviously, the actual DGLAP asymptotics of Eq. (13) is of the Regge type, it differs a lot from the conventional DGLAP asymptotics of Eq. (3) and looks similar to our asymptotics given by Eq. (10): incorporating the singular factors into DGLAP fits ensures the steep rise of g_1^{DGLAP} at small x and thereby leads to the success of DGLAP at small x . Ref. [9] demonstrates that without the singular factor $x^{-\alpha}$ in the fit of Eq. (11), DGLAP would not be able to operate successfully at $x \leq 0.05$. In other words, the singular factors in DGLAP fits mimic the total resummation of logarithms of x of Eqs. (4,10). Although both (13) and (10) predict the Regge asymptotics for g_1 , there is a certain difference between them: Eq. (13) predicts that the intercept of g_1^{NS} should be $\alpha = 0.57$. As α is greater than the non-singlet intercept $\Delta_{NS} = 0.42$, the non-singlet g_1^{DGLAP} grows, when $x \rightarrow 0$, faster than our predictions. However, such a rise is too steep. It contradicts the results obtained in Refs. [7] and confirmed in Refs. [8]. Usually, the DGLAP equations for the non-singlets are written in the x -space as convolutions of splitting functions P_{qq} with evolved parton distributions Δq and the latter are written as another convolution:

$$\Delta q(x) = C_q(x, y) \otimes \delta q(y), \quad (14)$$

with C_q being the coefficient function. Written in this way, Δq is sometimes believed to be less singular than δq because of the evolution. However applying the Mellin transform to Eq. (14) immediately disproves it.

IV. CONCLUSION

Comparison of Eqs. (3) and (13) shows explicitly that the singular factor $x^{-\alpha}$ in the Eq. (11) for the initial quark density converts the exponential DGLAP-asymptotics into the Regge one. On the other hand, comparison of Eqs. (10) and (13) demonstrates that the singular factors in the DGLAP fits mimic the total resummation of logarithms of x . These factors can be dropped when the total resummation of logarithms of x performed in Ref. [7] is taken into account. The remaining, regular x -terms of the DGLAP fits (the terms in squared brackets in Eq. (11)) can obviously be simplified or even dropped at small x so that the rather complicated DGLAP fits can be replaced by constants. It immediately leads to an interesting conclusion: the DGLAP fits for δq have been commonly believed to represent non-perturbative QCD effects but they actually mimic the contributions of the perturbative QCD, so the whole impact of the non-perturbative QCD on g_1 at small x is not large and can be approximated by normalization constants.

V. ACKNOWLEDGEMENT

B.I. Ermolaev is grateful to the Organizing committee of the workshop Spin-05 for financial support of his participation in the workshop.

-
- [1] G. Altarelli and G. Parisi, Nucl. Phys.B126 (1977) 297; V.N. Gribov and L.N. Lipatov, Sov. J. Nucl. Phys. 15 (1972) 438; L.N.Lipatov, Sov. J. Nucl. Phys. 20 (1972) 95; Yu.L. Dokshitzer, Sov. Phys. JETP 46 (1977) 641.
 - [2] B.I. Ermolaev, S.I. Manaenkov and M.G. Ryskin. Z.Phys.C69(1996)259; J. Bartels, B.I. Ermolaev and M.G. Ryskin. Z.Phys.C 70(1996)273; J. Bartels, B.I. Ermolaev and M.G. Ryskin. Z.Phys.C 72(1996)627.
 - [3] J. Blumlein, A. Vogt. Acta Phys. Polon. B27 (1996) 1309. J. Blumlein, S. Riemersma, A. Vogt. Nucl.Phys.Proc.Suppl.51C (1996) 30; Acta Phys. Polon. B28 (1997) 577.
 - [4] J. Kwiecinski and B. Ziaja. Phys. Rev. D60(1999)054004; J. Kwiecinski, M. Maul. Phys.Rev.D67(2003)03401; B. Ziaja. Phys.Rev.D66(2002)114017; A. Kotlorz, D. Kotlorz. Acta Phys.Polon. B34(2003)2943; Acta Phys.Polon. B35(2004)705.
 - [5] B.I. Ermolaev, M. Greco and S.I. Troyan. Phys.Lett.B 522(2001)57.
 - [6] D. Kotlorz A. Kotlorz. Acta Phys. Polon. B35(2004)2503; hep-ph/0407040.
 - [7] B.I. Ermolaev, M. Greco and S.I. Troyan. Nucl.Phys.B 594 (2001)71; ibid 571(2000)137; Phys.Lett.B579(321),2004.
 - [8] J. Soffer and O.V. Teryaev. Phys. Rev.56(1997)1549; A.L. Kataev, G. Parente, A.V. Sidorov. Phys.Part.Nucl 34(2003)20; Nucl.Phys.A666(2000)184; A.V. Kotikov, A.V. Lipatov, G. Parente, N.P. Zotov. Eur.Phys.J.C26(2002)51; V.G. Krivohijine, A.V. Kotikov, hep-ph/0108224; A.V. Kotikov, D.V. Peshekhonov hep-ph/0110229; N.I. Kochelev, K. Lipka, W.D. Novak, A.V. Vinnikov. Phys. Rev. D67(2003) 074014.
 - [9] B.I. Ermolaev, M. Greco and S.I. Troyan. Phys. Lett. B622 (2005) 93 (hep-ph/0503019).
 - [10] G. Altarelli, R.D. Ball, S. Forte and G. Ridolfi, Nucl. Phys. B496 (1997) 337; Acta Phys. Polon. B29(1998)1145;
 - [11] A. Vogt. hep-ph/0408244.